

DIFFRACTIVE PHOTOPRODUCTION OF MESON PAIRS IN THE SÖDING MODEL

N.I. Starkov

P.N. Lebedev Physical Institute, Russian Academy Sciences,
Moscow 117924, Russia

N.P. Zotov

D.V. Skobeltsyn Institute of Nuclear Physics,
M.V. Lomonosov Moscow State University, Moscow 119899, Russia

Abstract

We consider the diffractive photoproduction of $\pi^+\pi^-$, K^+K^- , D^+D^- meson pairs with high masses and transverse momenta in the framework of the Söding model. Total and differential cross sections are calculated for these processes. The role of form factors is stressed.

1 Introduction

The experimental results obtained by H1 and ZEUS Collaborations on HERA ep-collider for diffractive ρ^0 , ϕ , and J/Ψ meson production by real and virtual photons attract abnormally high attention [1]. The main reason of this attention is explained by the fact that transition from small virtuality photon ($Q^2 \simeq 0$) to deep inelastic scattering ($Q^2 > 4 \text{ GeV}^2$) connects with dynamic of a transition from soft diffractive γp interaction to hard one which is described by the vacuum number exchange, the so called Pomeron [2].

Recently H1 and ZEUS Collaborations published experimental data on diffractive ρ^0 meson photoproduction [3, 4]. These results confirmed the general properties of soft diffractive hadron processes which are well known, but also showed the necessity for further experimental and theoretical studies.

It is known that the cross section for diffractive dissociation of an incoming particle (a photon on our case) into low mass states contains at all three different contributions (see for instance [5]): 1) the term describing the direct production of hadron states (resonances) which then decay into final state hadrons, 2) the Drell–Hiida–Deck (DHD) nonresonant background term (more precisely the resonance like one) [6], and 3) the terms originating from final state interactions [7].

The DHD mechanism for the nonresonant background in ρ^0 meson photoproduction has been used by Söding [8]. The interference between the amplitude of DHD nonresonant $\gamma p \rightarrow \pi^+\pi^-p$ production and the amplitude of resonant $\gamma p \rightarrow \rho^0 p \rightarrow \pi^+\pi^-p$ production leads to a shift of ρ the mass spectrum. In general the correct account of the interference pion pairs produced by the DHD nonresonant mechanism and via the ρ^0 meson decay is a theoretical as well as an experimental problems [5, 7].

Therefore in this note we propose a simple model for nonresonant diffractive meson pair ($\pi^+\pi^-$, K^+K^- , D^+D^-) production which contains no diffractive resonance (ρ^0 , ϕ , ...) production.

The paper is organized as follows. In Section 2 we formulate the model (similar to the Söding model) for the meson pair photoproduction. In Section 3 we present our numerical results for diffractive photoproduction of meson pairs. Section 4 devotes qualitative study of the consequences of the absorption in the final state on t distribution of meson pairs. In Conclusions we emphasize an importance of the proposed model for a description of many background processes in a number of experiments on HERA collider.

2 A model for diffractive photoproduction of meson pairs

We use Söding model [8] for diffractive nonresonant photoproduction of meson pairs: $\pi^+\pi^-$, K^+K^- , and D^+D^- . In according with this model a photoproduction of meson pairs is described by two diagrams containing the exchange of a Pomeron (Fig.1), which correspond to the following formula for the differential cross section for meson pair production by transversely polarized photons ($Q^2 = 0$):

$$\frac{d\sigma^T}{dt} = \frac{\alpha}{2\pi^2} e^{Bt} \frac{\sigma_{tot}^2}{16\pi} \int \frac{z(1-z)dz d^2q_{t(-)}}{m_{(+)t}^2 m_{(-)t}^2} A F^2(q_{t(-)}^2), \quad (1)$$

where σ_{tot} and B are the usual parameters describing meson-proton scattering: the total cross section and the slope parameter of diffractive cone. The two-component transverse momentum of a meson $\vec{q}_{t(-)}$ is defined in the γp c.m.s. using the Sudakov decomposition [9] for 4-momentum of the negative charge meson in final state: $q_{(-)} = zQ + \beta p + q_{t(-)}$, with $z = (q_{(-)}p)/(Qp)$ and $\beta = m_{(-)t}^2/[z2(Qp)]$. Here Q and p are the 4-momenta of the photon and the proton (with $m_p = 0$), respectively. The factor A in (1) has the following form¹:

$$A = \left[\frac{(1-z)}{z} \frac{m_{(-)t}^2}{m_{(+)t}^2} q_{t(+)}^2 + \frac{z}{(1-z)} \frac{m_{(+)t}^2}{m_{(-)t}^2} q_{t(-)}^2 - 2\vec{q}_{t(+)}\vec{q}_{t(-)} \right] \quad (2)$$

and is equal to $\vec{q}_{t(-)}^2/[z(1-z)]$ at $t = 0$. The remaining kinematic variables in (1) and (2) have the following form: $m_{(-)t}^2 = \mu^2 + \vec{q}_{t(-)}^2$ is transverse mass of a meson, $t \simeq -(\vec{p}_t')^2$ and $\vec{q}_{t(+)} = -\vec{p}_t' - \vec{q}_{t(-)}$.

In (1) the functions $F(q_{t(-)}^2)$ are "form factors" of the two meson states connected by off mass shell mesons.² The exact expressions for these form factors are unknown. They can be written, for instance, in pole forms, where they are expressed through $\vec{q}_{t(-)}^2$ together with masses of suitable meson states. For π , K , and D mesons these masses can be chosen to be the masses of ρ , ϕ , and J/Ψ mesons (in the following named "version 1") or of $\pi(1300)$, ϕ , and J/Ψ mesons [10] ("version 2"). We use here both possibilities. As an example, for "version 2" the form factors for π pair photoproduction is

$$F_\pi(q_{t(-)}^2) = \frac{m_{\pi(1300)}^2}{m_{\pi(1300)}^2 + \vec{q}_{t(-)}^2}. \quad (3)$$

In the following, the total cross sections for πp and Kp scatterings in (1) are assumed to be described by the Donnachie - Landshoff parametrizations [11], which give at HERA energies ($\sqrt{s} = 100 GeV$) the following values: $\sigma_{tot}(\pi p) = 29.2 mb$, $\sigma_{tot}(Kp) = 25.1 mb$. For $\sigma_{tot}(Dp)$ we used the different values ranging from $3 mb$ ("version 1") up to $\sigma_{tot}(Dp) = \sigma_{tot}(\pi p)/2$ ("version 2"). The values of B - parameters in (1) are equal: $B_{\pi p} = B_{Kp} = 10 GeV^{-2}$ and $B_{Dp} = 6 GeV^{-2}$.

We would like to point out that the values of the cross section $\sigma_{tot}(Dp)$ and the slope parameter B_{Dp} are not determined enough exactly so far and the obtained results for the D meson photoproduction cross sections depend essentially on these ones and corresponding form factor.

¹ The approximate expression for A initially was obtained by M.Ryskin [10]

²In the original Söding model [8] these form factors are absent.

The distribution $d\sigma^T/dM$ over the mass of a meson pair, $M = (q_{(+)} + q_{(-)})$, can be obtained through the transition from z to M in eq. (1). We used the following relations valid at $t = 0$:

$$z(1-z)M^2 = \mu^2 + \vec{q}_{t(-)}^2, \quad dz/dM = 2Mm_t^2/[M^4(1-4m_t^2/M^2)] \quad (4)$$

3 Numerical results

The differential cross section $d\sigma/dt$ for the processes of diffractive photoproduction of meson pairs ($\pi^+\pi^-$, K^+K^- , and D^+D^-) are presented in Fig.2. The slopes of these cross sections are determined in the main by B parameters in (1).

We evaluated the values of these differential cross sections at $t = 0$. Based on the form factors of "version 1" we obtained $45.3 \mu b/GeV^2$ ($\pi^+\pi^-$ pairs), $14.5 \mu b/GeV^2$ (K^+K^- pairs) and $0.14 \mu b/GeV^2$ (D^+D^- pairs). "Version 2" resulted in $90 \mu b/GeV^2$, $14.5 \mu b/GeV^2$ and $3.5 \mu b/GeV^2$ correspondingly.

The transverse momentum distributions of outgoing mesons $d\sigma/dtdq_{t(-)}^2$ for the process $\gamma p \rightarrow K^+K^-p$ at a fixed values $|t| = 0$ and $2 (GeV/c)^2$ are shown in Fig. 3. These distributions at $|t| = 0$ and/or $|t| \neq 0$ but at high $|\vec{q}_{t(-)}| > 3 GeV/c$ have a behaviour $\sim |\vec{q}_{t(-)}|^{-6}$.

The cross sections $\sigma(|\vec{q}_{t(-)}| > |\vec{q}_{t(-)}|_{min})$ for $\gamma p \rightarrow \pi^+\pi^-p$, $\rightarrow K^+K^-p$, and $\rightarrow D^+D^-p$ processes shown in Fig.4. The total cross section for diffractive meson pair photoproduction was obtained by integrating expression (1). Using the form factors of "version 1" results in $4.6 \mu b$ for $\pi^+\pi^-$ pairs, $1.4 \mu b$ for K^+K^- pairs and $\sim 0.02 \mu b$ for D^+D^- pairs. Based on the form factors of "version 2" we obtained: $10 \mu b$, $1.4 \mu b$ and $\sim 0.5 \mu b$ correspondingly.

The results for the mass distributions of diffractive photoproduction of pion and kaon pairs are shown in Fig.5. The curves (1) and (2) for the pion pair photoproduction correspond to two versions of the form factor (3). The value of the background in the ρ^0 mass region $0.55 < M < 1 GeV$ somewhat larger than the ZEUS estimates[4] and at larger mass there is a strong dependence on the choice of the form factor.

The total cross sections obtained by integrating $d\sigma/dM$ over the mass M of a meson pairs are presented in Fig.6. The values of these cross sections at high M coincide approximately with those obtained by integrating expression (1) over z and $q_{t(-)}$.

As we have pointed in Introduction meson pair photoproduction cross sections contain final state interaction contribution besides resonant and non-resonant ones. The effects of rescattering in final state (or final state absorption) for $\pi^+\pi^-$ - photoproduction can lead to 20-40% corrections [5, 12]. But for photoproduction of heavy mesons (K and D) our model is more adequate.

4 Some remarks on absorptive corrections

The expression for differential cross section (1) takes into account the contribution of the elastic meson - proton scattering and also all the rescatterings of one meson on proton (see Fig.1). Then for the elastic (diffractive) photoproduction of meson pairs one should include two kinds of the absorptive corrections in the final state: the elastic rescattering of mesons on each other and one of both mesons off the target (proton). The former corrections are essential for the resonance (for instance ρ meson) production of meson pairs can be take into account by the special prescription [7, 12] and do not affect considerably on the slope parameter of differential cross section.

The things go quite differently for the absorptive corrections connected with the rescattering of two meson system as whole off the target. These corrections lead to strong effects for t -dependence of meson pair differential cross section (see, for instance [5]).

Here we restrict ourselves to the allowance of these corrections on the phenomenological level, when in the eikonal approximation the dependence $d\sigma/dt \sim \exp Bt$ is replaced by [12]:

$$\frac{d\sigma}{dt} \sim \left[\exp(Bt/2) - \frac{\sigma}{16\pi B} \exp(Bt/4) \right]^2 \quad (5)$$

The absorptive corrections in final state may be enhanced due to inelastic diffractive intermediate states in the proton vertex of diagram in Fig. 1, then σ is replaced by $\lambda\sigma$ [5, 13] in the second term of eq. (5). The phenomenological parameter λ should have the following value: $\lambda \approx 1 + \sigma_D/\sigma_{el}$ [14]. Some estimates for NN -interactions give $\lambda \approx 1.2$ [15]; in the region of the laboratory energy $E_L = 10 - 10^3 \text{ GeV}/c$ the ratio $\sigma_D/\sigma_{el} \approx 1$ and $\lambda \approx 2$. Therefore in our calculations we used the following values: $1.2 < \lambda < 2$ [5].

The results of our calculations of differential cross section of the process $\gamma p \rightarrow \pi^+\pi^-p$ by the formula (5) are illustrated in Fig. 7 at suitable set of parameters (for arbitrary normalization of $d\sigma/dt$): $\sigma = 28 \text{ mb}$, $B = 7 (\text{GeV}/c)^{-2}$ and $\lambda = 1.3$.

In rough approximation for the allowance of the absorptive corrections differential cross section $d\sigma/dt$ has a dip at $t \simeq 0.75 (\text{GeV}/c)^2$, which may be filled in a more realistic model. But the main point is that the $d\sigma/dt$ changes of the slope parameter at high t (at $|t| > 0.8 (\text{GeV}/c)^2$), which is observable in the preliminary ZEUS experimental data for $\gamma p \rightarrow p\pi^+\pi^-$ at $M_{\pi^+\pi^-} = 0.5 - 1.1 \text{ GeV}$ in the region $W = 64 - 90 \text{ GeV}$ and $0.25 < Q^2 < 0.8 \text{ GeV}^2$ [16]. Similar dips may appear in differential cross sections of diffractive J/ψ and ρ^0 -productions in DIS [17, 18].

5 Conclusions

We presented a new model for diffractive meson pair photoproduction which we applied to estimate the non-resonant meson pair photoproduction cross sections. We showed that the results strongly depend on the choice of form factors. The t -dependences of differential cross sections are determined essentially by the absorptive corrections in final state. Although the non-resonant cross-sections are small in comparison with the resonant diffractive meson pair photoproduction, they strongly affect background processes in ρ^0 , ϕ and J/ψ meson production by real and virtual photons.

First of all the known mass shift effect for ρ - meson photoproduction which is controlled [8] by the Söding mechanism of the non-resonant $\pi^+\pi^-$ pair photoproduction [3, 4]. The Söding mechanism contribution is visible at low $Q^2 \neq 0$ ³ and it can be estimate in the framework of our model (see also [19]).

The second our estimations for non-resonant diffractive K^+K^- meson photoproduction contribution are important for correct extraction of K^- meson photoproduction cross section at HERA energies. As was noted by Pumplin [12] the process $\gamma p \rightarrow K^+K^-p$ is interesting because it could be studied at low K^+K^- masses without interference from ϕ -resonance production.

Our proposed D^+ meson photoproduction mechanism should give contribution to the μ - production at large transverse momentum q_t . These muons can give large background for muon pairs from J/ψ meson production on HERA collider.

Finally the formula (1) can be used for extraction of the elastic $\pi^\pm p$, $K^\pm p$ and $D^\pm p$ cross sections [19]⁴ if we have experimental values for non-resonant diffractive $\pi^+\pi^-$, K^+K^- and D^+D^- pair photoproduction cross sections.

³We are grateful to H. Beier for this remark.

⁴ Initially it was proposed by M. Arneodo

6 Acknowledgements

One of us (N.Z.) would like to thank M. Ryskin for the original suggestion to study diffractive non-resonant meson pair photoproduction and for useful discussions, A. Proskuryakov for discussions of correlations of the resonant and background contributions in meson pair photoproduction and J. Repond for the careful reading of the manuscript and useful remarks.

References

- [1] L.P.A. Haakman, A. Kaidalov, J.H. Koch, Phys. Lett. B365 (1996) 411.
E. Gotsman, E.M. Levin, U. Maor, Nucl. Phys. B464 (1996) 251.
L. Frankfurt, W. Koepf, M. Strikman, Phys. Rev. D54 (1996) 3194.
J. Nemchik, N.N. Nikolaev, E. Predazzi, B.G. Zakharov, Phys. Lett. B374 (1996) 199.
- [2] J. Nemchik, N.N. Nikolaev, B.G. Zakharov, Phys. Lett. B341 (1994) 228.
H. Abramowicz, L. Frankfurt, M. Strikman, preprint DESY 95-047.
A. Levy, preprint DESY 95-204.
- [3] H1 Coll., S. Aid et al., Nucl. Phys. B463 (1996) 3.
- [4] ZEUS Coll., M. Derrick et al., Z. Phys. C69 (1996) 39.
- [5] N.P. Zotov, V.A. Tsarev, Sov. J. Part. and Nucl. 9 (1978) 266.
- [6] S.D. Drell, K. Hiida, Phys. Rev. Lett. 7 (1961) 199.
R. Deck, Phys. Rev. Lett. 13 (1964) 169.
- [7] M. Ross, L. Stodolsky. Phys. Rev. 149 (1966) 1172.
- [8] P. Söding, Phys. Lett. 19 (1966) 702.
- [9] V.V. Sudakov, ZhETF 30 (1956) 87.
- [10] M.G. Ryskin, private communication.
- [11] A. Donnachie, P.V. Landshoff, Phys. Lett. B296 (1992) 227.
- [12] J. Pumplin, Phys. Rev. D2 (1970) 1859.
- [13] M. Ross, F.S. Henyey, G.L. Kane, Nucl. Phys. B23 (1970) 269.
G.L. Kane, A. Seidl, Rev. Mod. Phys. 48 (1976) 309.
- [14] K.G. Boreskov, A.M. Lapidus, S.T. Sukhorukov, K.A. Ter-Matrtirosyan,
Yad. Fiz. 14 (1971) 814.
- [15] M. Uehara, Progr. Theor. Phys. 56 (1976) 1183.
- [16] T. Doeker, private communication.
- [17] L.L. Jenkovszky, E.S. Martynov, F. Paccanoni, hep-ph/9608384.
- [18] E. Gotsman, E. Levin, U. Moar, hep-ph/9701236.
- [19] M.G. Ryskin, Yu.M. Shabelski, hep-ph/9701407.

Figure captions

1. Diagrams for diffractive meson pair photoproduction in Söding model.
2. The differential cross sections $d\sigma/dt$ for diffractive meson pair photoproduction.
3. The transverse momentum distributions for $\gamma p \rightarrow K^+ K^- p$ at fixed values of $|t|$.
4. The integral cross sections for diffractive meson pair photoproduction processes as a function of q_{min} .
5. The mass distribution $d\sigma/dM$ over the mass of two π^- and K^- -mesons: the curves (1) and (2) for the version 1 and 2 of form factors.
6. The total cross sections $\sigma(M_{max})$ versus the mass of two π^- and K^- -mesons: curves as in Fig.5.
7. The differential cross section for the process $\gamma p \rightarrow \pi^+ \pi^- p$ obtained by the formula (5).